

Council of Europe

CONVENTION ON THE CONSERVATION OF EUROPEAN WILDLIFE
AND NATURAL HABITATS

Action plan for *Margaritifera margaritifera* in Europe

Authors

Dr. Rafael ARAUJO and Dr. M^a Ángeles RAMOS
Museo Nacional de Ciencias Naturales (CSIC)
José Gutiérrez Abascal, 2. 28006-Madrid, Spain
Tel 34-914111328, Fax 34 915645078
e-mail rafael@mncn.csic.es, m.ramos@mncn.csic.es

This report has been drafted with the support of the Government of Belgium - Ministère de la région wallone

CONTENTS

SUMMARY	5
1. INTRODUCTION	6
2. BACKGROUND INFORMATION	7
2.1. Systematics	7
2.2. Description of the species	7
2.3. Distribution and population numbers	8
2.3.1. World distribution	8
2.3.2. European distribution by countries	8
2.4. Life history	13
2.4.1. Life cycle	13
2.4.2. Habitat requirements	14
2.5. Threats and limiting factors	14
2.5.1. Habitat loss and habitat alteration	15
2.5.2. Human activities	16
2.6. Conservation status and current legal protection	16
2.6.1. International agreements	16
2.6.2. Listing of individual countries	17
3. GOALS AND OBJECTIVES	25
3.1. Policy and legislation	25
3.2. Species and habitat protection	26
3.2.1. Develop and implement a management plan to prevent further decline	26
3.2.2. Conduct population surveys	26
3.3. Conflicts with humans	27
3.4. Public awareness	27
3.5. Population conservation	27
3.6. Artificial reproduction	28

3.7.	Artificial propagation and reintroduction	28
3.8.	Natural recovery	28
3.9.	Monitoring and research	28
4.	CONSERVATION ACTIONS	29
4.1.	International actions	29
4.2.	Listing of countries with necessary actions	29
5.	REFERENCES	36
6.	ACKNOWLEDGEMENTS	39
7.	APPENDIX 1: Summary on protection and management measures on <i>M. margaritifera</i>	40
8.	APPENDIX 2: List of contributors to this report	41

SUMMARY

The freshwater pearl mussel *Margaritifera margaritifera* (L.) was probably the most abundant bivalve in ancient rivers all around the world. Although it is very threatened, populations still exist in many European countries. The species is protected under the Habitats Directive (Appendix II and Appendix V) and the Bern Convention (Appendix III). In the IUCN 1996 Red Data Book it is listed as Endangered.

The species needs a fish of the family Salmonidae to host the metamorphosis of its larval stage, called glochidium. Thus, alterations to the habitat of the host fish also work against the mussel. The river bed substratum is of great importance to the mussels, and determines the river areas where they can survive. Clean gravel and sand are essential for a healthy population, being very important for juveniles because if the substratum becomes clogged with silt, oxygen can no longer reach them and they die. For them to thrive, there needs to be enough brown trout or salmon with viable natural reproduction. Anomalies in feeding conditions within mussel habitats and in their environments are the clues to the rarification of this species.

The causes of the species decline are not completely known, but habitat alteration and fragmentation are probably the origin. Taking into account the data provided by each country's National Nature Conservation Agencies, the main reasons for the decline in *M. margaritifera* are anthropogenic influences on aquatic systems: eutrophication, impoundment, river regulation, drainage, sewage disposal, dredging, farms, new agricultural land use, loss of forests and natural river banks, water pollution, acidification, pesticides, and introduced exotic fish species. Factors influencing the abundance of fish hosts are also responsible. Data suggest that around 90% of the European specimens of *M. margaritifera* disappeared in the twentieth century. Overfishing for pearls was one of the main factors, particularly in some countries.

Legislation, rational water use, research, management (including species reintroduction), monitoring and education/consultation will be of foremost importance for species conservation. As a freshwater animal, *M. margaritifera* comes into constant conflict with people, mainly due to the large-scale engineering works for water exploitation (canalisation, dredging, regulation, impoundment, power plants, etc.). Due to this fact, the first thing is to establish the priorities in the respective countries. There is no sense in demanding freshwater species conservation at the same time as main roads, dams and macro-changes in the landscape make the freshwater ecosystem unsuitable for those species. As fertility is maintained even in sparse populations and in pollute rivers, the populations should recover if the causes of decline are removed.

Further studies are needed to fill the gaps in the known distribution of the species in Europe.

Protection of European freshwater systems harbouring populations of *M. margaritifera* under the Natura 2000 Network or as UNESCO Biosphere Reserves may be essential for species survival.

1. INTRODUCTION

The freshwater pearl mussel *Margaritifera margaritifera* (L.) was probably the most abundant bivalve in ancient oligotrophic rivers all over the Northern Hemisphere (Holarctic Region). Although dense populations have been described from Northern Europe, *M. margaritifera* lived on both sides of the Atlantic. The identity of similar species living on the North American West coast (*M. falcata*) and in the Far East (*M. laevis*) has not yet been clarified. The best preserved populations are at present in Western Russia in rivers on the Kola Peninsula.

Although in decline, populations still exist in many European countries. The species is protected under the Habitats Directive (Appendix II and Appendix V) and the Bern Convention (Appendix III). It is listed as Endangered in the IUCN 1996 Red Data Book. Some European countries harbouring populations of the species have developed action plans to protect it (see below); however, this document is the first attempt at a Pan-European Action Plan for the species.

Until 1985, the main bibliography about the species was summarised by Jungbluth, Coomans & Grohs (1985), but a lot of new information is currently flourishing (see references).

All freshwater mussels share a complex life cycle because they require a vertebrate host, usually a fish during their larval stage. This microscopic thin-shelled larva (glochidium) that the mussels brood and release by the millions normally has hooks and/or teeth to attach itself to the fish body where it encapsulates and spends several weeks completing its development until its recruitment as a metamorphosed benthic juvenile. This is the mussel's dispersal method. For this reason, maintaining natural populations of native fish is essential for the conservation of the naiads. Many data exist about the specificity between the glochidium of *M. margaritifera* and fish of the Salmonidae family, especially *Salmo salar* and *Salmo trutta fario*. Thus, alterations to the habitat of the fish hosts also work against the mussel.

Margaritifera margaritifera, as a freshwater animal, is in constant conflict with people, mainly as a result of big engineering works for water exploitation (canalisation, dredging, regulation, impoundment, power plants, etc.). When preparing this action plan, we tried not to obviate the "human factor" and to work with two kinds of approaches in accordance with Council of Europe recommendations (Council of Europe, 1996; Machado, 1997) on integrated ecosystem management and control of processes that negatively affect biodiversity.

In this dossier we have attempted to compile information from the countries with living populations in order to fulfil the necessary objectives required in drafting an action plan for *M. margaritifera*. These objectives are as follows:

1. To update the distribution and biological data.
2. To establish the threats hanging over the species.
3. To establish useful and realistic conservation measures.
4. To recover populations.

The information included in this report has been compiled from the bibliography, personal data and data provided by the different governmental agencies in countries where the species is thought to live. The answers to our questionnaire were very heterogeneous, as can be seen from this report, probably according to the level of knowledge in each country. Being aware of the risk of this heterogeneity, we

preferred to assume it, including detailed information from countries where it was available. The report contributors are listed in Appendix 1. We are very grateful to all of them for their help.

2. BACKGROUND INFORMATION

2.1. Systematics

Phylum Mollusca

Class Bivalvia

Order Unionoida

Family Margaritiferidae

Common name: Freshwater pearl mussel, Ostra de río, Almeja perlífera, Madreperla de río, Ameixa de río, Mulette perlière, Moule perlière, Flussperlmuschel, Elvemuslingen, Flodpärl musslan, Ziemelu upesperlene, Upes perlgliemene, Presnovodnaja zhemchuznitsa.

The genus *Margaritifera* was described by Schumacher in 1816, with *Mya margaritifera* Linnaeus, 1758 (= *Margaritifera margaritifera*) as the type species. However, its taxonomy is still unclear and, therefore, also the number of living species it contains. Although it is a declining genus, with all its species endangered, living species of *Margaritifera* have been recorded from North America, Europe and Asia, with only two species in Europe, *M. margaritifera* and *M. auricularia*.

Margaritifera, with the North American genus *Cumberlandia* and the uncertain Asiatic genus *Margaritanopsis*, constitute the family *Margaritiferidae*, including the oldest species of the Unionacea. It includes species with characteristic conchological and anatomical features (i.e. absence of diaphragm, incubation in the four gills, marsupium without water tubes, hookless glochidium) different to other naiads (families Unionidae, Mutelidae, Etheridae) (Davis & Fuller, 1981; Smith & Wall, 1984). These differences have recently been supported using molecular data (Davis & Fuller, 1981; Lydeard, Mulvey & Davis, 1996).

The taxonomic status of the Irish hardwater *M. durrovensis* is scientifically discussed in relation to *M. margaritifera* (Phillips 1928; Chesney, Oliver & Davis, 1993; Moorkens & Costello 1994). At present, only *M. margaritifera* and *M. auricularia* appear on the list of wildlife species under the Council of Europe's Convention on the Conservation of European Wildlife and Natural Habitats (Bern Convention 1979). However, the European Union Directive on the Conservation of Natural and Semi-Natural Habitats and of Wild Fauna and Flora (Directive 92/43/EEC, Habitats Directive), recently listed *M. durrovensis* as *M. durrovensis* (*M. margaritifera*) in Annex II and Annex V (Directive 97/62/CE). In the International Union for the Conservation of Nature and Natural Resources Red Data List (IUCN 1996) *M. durrovensis* is listed as Critically Endangered.

2.2. Description of the species

Adults present a large, dark, fragile and elongated shell up to 15 cm long. The umbones do not rise above the shell and are generally very eroded. The longer region of the shell contains the posterior part of the body, including the water interchange inhalant and exhalant apertures; the foot aperture is located in the anterior region.

The inner side of the shell is pearl white, sometimes with iridescent colours. The pallial line is less obvious than in *M. auricularia* and the inner muscle scars are obvious, mainly the anterior one. The

hinge has vestigial posterior lateral teeth, practically inexistent, and pseudocardinal teeth less developed than in *M. auricularia*, one in the right valve and two in the left, the posterior one being less developed.

The foot is white and very big, capable of becoming as large as the whole shell when protruding. There are no real siphons, as only a transversal thickening on the inner side of the posterior mantle (the diaphragm) maintain the exhalant and inhalant apertures separate in live specimens. The edge of the mantle in this area is thickened by rows of papillae. Anteriorly, the labial palps, two on each side, move the water currents in order to select the food which will be ingested through the mouth, between the palps, once it has been sorted by the gills. The latter are under the mantle, two inner and two external, all without water tubes, an important difference with other families of Unionacea. In the dorsal part of the foot and embedded in the visceral mass is the gonad, which is sometimes hermaphroditic.

The glochidia are very small (60 x 70 µm). They were recently described as having neither spines nor teeth (Nezlin *et al.*, 1994; Pekkarinen and Valovirta, 1996) although Harms (1907, 1909) described very minute teeth at the ventral border in the first description of this larva.

2.3. Distribution and population numbers

2.3.1. World distribution

Margaritifera margaritifera lives on both sides of the Atlantic in clean streams and rivers with healthy populations of salmonids throughout much of the Holarctic region. It is known from north-west Russia, Northern Europe, eastern North America and Canada. In Europe, it occurs in rivers of the Arctic and Atlantic oceans. Its range is bounded by the Alps to the South and by the basin of the Severnaya Dvina River to the East (Ziuganov *et al.*, 1994). In North America, it occurs on the Atlantic coast from Newfoundland, Canada down to Delaware and Pennsylvania. To the West, the range is bounded by the Appalachians.

Population numbers are declining in all countries. The species is nearly extinct in many areas and only undisturbed river basins, such as those on the Kola Peninsula, maintain flourishing populations with important juvenile recruitment.

2.3.2. European distribution by countries

- Austria

The species is endangered and has been reduced by about 97-98% since the late 19th century. Of the 231 river sections sampled, 27 populations were located (Moog *et al.*, 1998). Most of the remaining populations are either senescent or declining (Moog *et al.*, 1998). There is an estimated total of 50,000 individuals covering approximately 2,520 km². The largest population so far discovered contains about 20,000 individuals, and 19 populations have less than 1,000 (Moog *et al.*, 1998).

- Belgium

The past distribution of the species included only south-eastern rivers all of them flowing in the Ardenne region: Amblève, Ourthe, Lomme, Lesse rivers (Adam, 1960). It was also known from the

Schwalm (Jungbluth, 1993) and the River Our, mainly in the Luxembourg section (Trois Frontières, Frein, Moulin de Kalborn, Tintesmühle, Grossenaul) (Birringer & Truffner, 1990). At present its distribution is poorly known, and although the data are contradictory it seems that only a few aged specimens have been located in small streams of the Rhine catchment (including the river Our) and from rivers belonging to the basin of some important tributaries of the Meuse (Vrignaud, 2000; Roland Libois, pers. obs.).

- Czech Republic

There are about 130,000 specimens in 15 localities of different rivers and brooks. The main population is in the River Blanice, with 115,000 individuals (Hruska, 1999).

- Denmark

The species is only found along a relatively short stretch of the River Varder in the north-west of Jütland, the last records being from 1974. There is no current data on the species in the river, and there is a need to investigate if the species still occurs in the locality and if possible specimens can still breed (Stoltze & Pihl, 1998).

- Estonia

According to data from 1998, there is only one locality with an old population of around 40,000 specimens.

- Finland

At the beginning of this century there were approximately 200 rivers with freshwater pearl mussel populations. Today, only 25% of these rivers still contain populations. In southern Finland, the populations have declined radically, only in one of the eight rivers is there a breeding population (Valovirta & Yrjänä, 1996). The species never occurred in the lake district of Central Finland. The last estimation (1999) yielded about 3 million individuals, 90% of them in eastern Lapland (65°N-69°N). Together with its tributaries, the River Lutto harbours the most numerous populations. The current total is estimated at least 1.5 million.

- France

Historically the species was present in water courses of the oldest mountains such as the Armorican Massif, the Vosges, the Ardennes, the Central Massif and the western Pyrenees. The cites of the Alps and the Jura are probably erroneous (Vrignaud, 2000). The population has decreased by about 90% in the last century, the estimated number of individuals being about 100,000. The main decrease occurred in Brittany, Normandy, Vosges and the Massif Central (Cochet, 1999).

Data derived from surveys since 1993 report the species from at least 74 rivers in 26 *départements* as follows (Guilbot *et al.*, 1998): 14 Armorican Massif, 57 Central Massif and Morvan, 1 Vosges and 2 Pyrenees. News of breeding populations only comes from rivers in Morvan, the Central Massif and the Pyrenees.

- Germany

The information on *M. margaritifera* detailed by regions is as follows:

- Schleswig-Holstein: No viable populations have ever existed.
- Niedersachsen: A population of 2,000-2,500 specimens exists in the Lachte-Lutter stream system.
- Nordrhein-Westfalen: The species is extinct. The last small populations of the Eifel mountain system were translocated four years ago to Belgium.
- Rheinland-Pfalz: Although formerly occurring in several rivers of this state the species is now regarded as extinct, except for the population in the river Our, shared with the country of Luxemburg.
- Hessen: No data on presence of the species are available. There exists one relic population of about a dozen of aged specimens in the Vogelsberg mountain (Kremer, pers. comm.).
- Baden-Württemberg: Listed as threatened with extinction in the Red List of Endangered Snails and Mussels in Baden-Württemberg (1995), but regarded as extinct nowadays. The species does not occur in this state. Mussels were artificially introduced in the 18th century into the Steinach river system (Odenwald mountains), being the only successful introduction in this state, but the population is currently (1978) extinct.
- Bayern: In the nineteenth century, the population was estimated at 20 million individuals. There are now approximately 200,000 in 60 populations in freshwater systems of the Bayerischer Wald, Oberpfälzer Wald and Fichtelgebirge mountains. In the last decade, only 10 populations contained juveniles, but the reproductive rate is not enough to ensure their future survival.
- Thüringen: The species in this state only lived in small freshwater systems, but went extinct in the last or first half of the twentieth century. Two no-chance specimens exist in the Wettera River near Schleiz.
- Sachsen: In the Wolfsbach, the population in the watershed system of the Saale stream numbers 1,700; in the Triebelbach and Raunerbach in the watershed system of the Weisse Elster stream, there are approximately 100 specimens.

- Ireland

According to recent reports (Moorkens, 1999), the species is still widespread, with at least well-dispersed remnant populations around the country away from the central limestone plain in soft waters lying mainly on granite or sandstone bedrock, except for the *M. durrovensis* population in the lime-rich River Nore. Of 32 living populations, only eight have young mussels. In some of these populations, the last successful recruitment of young dates back to the 1960s or early 1970s (Moorkens, 1999). However, populations are numerous and apparently successfully reproducing in a small number of rivers. Recent survey (1999) demonstrates that two of these each contain in excess of 2,500,000 living, adult mussels (Speight, pers. comm.).

- Latvia

There are five populations in five different streams with total length of 25 km in the eastern part of the country. Population number (about 11,800 specimens) and size have decreased by more than 80% since the beginning of the 20th century. There are no specimens younger than 35-40 years (63 mm).

- Lithuania

The last record of the species is from 1990 in the east of the country.

- Luxembourg

Until the first half of the twentieth century the species was widely distributed in the Ardennes region, being common in Ösling streams (Ferrant, 1892, 1902). Altitude is between 180m (low section of the River Our) and 400m (east of the Ardennes). A few shells of recently dead mussels were found 30 years ago in the Rivers Troine, Clerve, Wiltz, Sure and Our (Groh, 1999).

West of the Rhine, the populations of the Vosges, the Eifel and the Hunsrück are extinct and in regression in the Ardenne. In this region, some rivers, either in Belgium (Amblève, Ourthe, Lomme, Lesse) or in Luxembourg (Troine, Cleve, Wiltz) where thriving populations were mentioned in the past, require further research to assess if live populations still exist there. Fortunately, the River Our and some small rivers of southern Belgium are still populated, sometimes by a reasonable number of mussels (>1000 ind) (Libois, pers. comm.). Therefore, these streams and the River Our seem to be the most important for the conservation of the species in the north-western continental Europe. However, the populations seem to be at risk because it seem that they contain only aged non-breeding specimens. In the River Our, the population was estimated at about 3,000 adults in 1989, while in 1999 only 1,500 specimens were estimated along a 20 km section of the river (Groh, 1999).

- Netherlands

The species does not occur in this country. There was one unsuccessful recolonization attempt in 1991.

- Norway

The populations of the species have shown a steady decrease over a long period. In many localities they have become extinct or recruitment has ceased, resulting in populations top-heavy with old individuals (Larsen, 1997). The species occurs in 340-350 watercourses in all the counties of Norway, but is most numerous along the coast. The greatest decline has been recorded in the counties of Vest-Agder, Aust-Agder and Rogaland. Little information is available for the rest of the country.

- Poland

The species has been extinct in Poland for fifty years. It was fairly abundant in Lower Silesia until the end of the Middle Ages. It was also known to be a common species in a few Sudetic streams in the early 1900s. Its occurrence in Poland has not been confirmed since the 1930s.

- Portugal

There are only old published reports of the species in Portugal (Nobre, 1941) and specimens stored at the Museu de Historia Natural (Faculdade de Ciencias do Porto), which were collected in the Douro, Sousa and Vouga rivers. A very recent record of the species is reported in the Rabasal River (Rolán, pers. comm.).

- Slovakia

The species did not originally occur in this country.

- Spain

Results point to the decline of Spain's salmon rivers during the twentieth century although new records of the mussel have been added to those reported by Bauer (1986), extending the species's distribution area to other river basins in Asturias and Zamora (Araujo & Ramos, in press; Ramos, 1998). The species is present in 14 rivers of the following provinces: Pontevedra, A Coruña, Lugo, Asturias and Zamora. The best populations occur in the Narcea River (Asturias), where densities are between 14.5 ± 15.9 ind/m² and 76 ± 5.72 ind/m² and the existence of juvenile recruitment has been demonstrated (INDUROT, 1999; Álvarez-Claudio *et al.*, 2000). New surveys are needed to ascertain the real distribution and densities of the species.

- Sweden

The freshwater pearl mussel was the subject of inventories in the field in the period 1980-1997 in 18 of the 21 counties, being absent in the other three. It was present in 370 of the 1100 watercourses surveyed. About three million individuals were estimated at 53 carefully studied sites, although following recent data from the WWF, the current number of specimens is about eight million (Henrikson, pers. com.). Compared with data from the first half of the twentieth century, the species has disappeared from more than 35% of its former sites, and recruitment fails at approximately 75% of sites. Nowadays, recruitment occurs at only one third of the inhabited sites. The healthiest populations are found in the northern and extreme south-eastern parts of Sweden. A common pattern is that populations in the main streams have disappeared, while isolated fragments or more widespread populations exist in the tributaries.

- United Kingdom.

— Scotland: Scotland is the UK stronghold for the freshwater pearl mussel. Historical records indicate that the freshwater pearl mussel was originally present throughout mainland Scotland (except in the River Tweed catchment and in the sandstone area of Caithness) and on some of the larger Western Isles, occupying over 150 rivers in total. All data suggest that in Scotland populations of the species are undergoing a catastrophic decline. In a recent survey, only 48 of the 148 sites that were known to be occupied 100 years ago retain functional populations; of these only 15 populations show evidence of recruitment, ten of them having more than 500 mussels per 50 m transect. The rate of extinction has doubled since 1970, with an average rate of loss of two populations per year since then (Cooksley, 1999).

— England: Few populations of the species remain in England. There are historical records from Devon, north Cornwall, the rivers Wye and Severn, the North Yorkshire Moors, the North Tyne and

Cumbria. A survey of seven of these rivers and one small stream found generally sparse populations, with only four having evidence of recruitment. Of these four rivers there is only one potentially viable population, numbering several hundred thousand adults, however, mortality greatly exceeds recruitment in this population. (Cooksley, 1999).

— Wales: In Wales the species is very scarce. There is historical evidence of large populations in the Rivers Conway and Wye, but which are now virtually extinct. There are records from streams in Pembroke, Carmarthenshire, Cardiganshire, Glamorganshire, Denbighshire, Snowdonia and some tributaries of the River Severn. The south-west and most of the Snowdonian populations now consist of only a few hundred senescent individuals. Only one Snowdonian population, which numbers approximately 3000 adults, shows evidence of some recent recruitment (Cooksley, 1999).

— Northern Ireland: Historically, populations of *M. margaritifera* were found in the Foyle, Bush and Erne catchments, and the central Loagh Neagh feeder streams. Recent surveys have found live mussels at only 20 sites (3 rivers) out of 250 surveyed. Juveniles are virtually absent in these populations and recruitment rates are well below those needed to sustain them (Cooksley, 1999).

2.4. Life history

2.4.1. Life cycle

All freshwater mussels share a complex life cycle, requiring a vertebrate host, usually a fish, during their parasitic larval stage. These microscopic thin-shelled larvae (glochidium), which are brooded and released by the millions, were presumed in the nineteenth century to be a parasitic species of fish (*Glochidium parasiticum*). The glochidium usually has hooks to attach itself to the fish body (fins or gills), where it becomes encapsulated for several weeks before changing into a free-living juvenile. The glochidium of *M. margaritifera* is of the hookless type, but presents very minute teeth at the ventral border (Harms, 1907, 1909) although this larva was recently described as having neither spines nor teeth (Nezlin *et al.*, 1994; Pekkarinen and Valovirta, 1996).

Information on the relationship between mussel and fish hosts is essential in any attempt to preserve endangered freshwater mussels. A lot of published information exists on the specificity between the *M. margaritifera* glochidium and salmonid fish, especially *Salmo salar* and *Salmo trutta fario*.

Once encysted in the fish gill filaments, the glochidium metamorphoses during a period of time which lasts depending on temperature. Two kinds of strategies have been described in the case of *M. margaritifera*, from 20-60 days to seven-nine months, although both strategies have been observed in the same population (Ziuganov *et al.*, 1994). Very little is known about juvenile habitat once they have emerged from the fish gill. Studies on this subject are currently underway in several European countries.

Sexual maturation occurs between seven and 15 years of age (Meyers & Milleman, 1977; Young & Williams, 1984) and the fertile period may be prolonged into old age. It is supposed to be a dioecious species although several cases of hermaphroditism have been reported (Bauer, 1987; Hanstén, Pekkarinen & Valovirta, 1997; Grande, Araujo & Ramos, 2001). According to Bauer (1987), the species can change its sex towards hermaphroditism under conditions of population stress. In all known European populations, females are gravid for two-three months from June, and glochidia are present in the fish gills from August.

The larvae are incubated in the four gills as in all other species of the family. An average of 9.8 million glochidia were found to be produced per female (Ross, 1992).

According to Bauer (1992), the species lifespan is between 30 and 132 years and the maximum shell length between 80 to 145 mm depending on the population (latitude and altitude). When there is enough available food, juveniles grow rapidly, but population lifespan is shorter. The largest known specimen is from the River Keret (Karelia, Russia) with 162 mm and the absolute theoretical maximum age is 167 years old (Ziuganov *et al.*, 1994).

Recent molecular data dealing with populations from Spain to Kola (Machordom *et al.*, in prep.) indicates that they can be considered as a metapopulation due the slight genetic differentiation among the specimens studied.

2.4.2. *Habitat requirements*

It was a common species in upper and middle river sections, mainly in rapid, clean and oligotrophic waters. *M. margaritifera* lives in granitic areas with low calcium rates. It lives partially burrowed in gravels and stones, sometimes behind stones or rocks, taking advantage of the available currents. Juvenile specimens (under 50 mm) can live completely burrowed (up to 20 cm) in the substratum. There is one hardwater population in Ireland known as *Margaritifera durrovensis* whose taxonomic status is under discussion (Chesney, Oliver & Chesney, 1993; Moorkens, 1996).

The river bed substratum is very important to mussels and determines the river areas where they can survive. Clean gravel and sand are essential to a healthy population (Moorkens, 1999) and it is very important for juveniles because if the substratum becomes clogged with silt, they can no longer obtain oxygen and so they die.

Margaritifera margaritifera lives in typical salmon rivers at depths of between 0.5 and 2 m (Ziuganov *et al.*, 1994), but it may also be common at greater depths. It can tolerate temperatures of up to 28°C for a short time (10-20 min) and is able to survive out of water for 30 days at 15°C. It avoids turbid waters and waters with low oxygen levels. Except for people, it has no natural enemies although it may be predated by the muskrat (*Ondatra zibethicus*). For it to thrive, there needs to be sufficient brown trout or salmon with viable natural reproduction.

The fact that old mussels can breed and that high mortality is not common suggests that juvenile interstitial phase may be the most sensitive period of the life cycle. Knowledge of auto-ecological requirements during the first 4-5 years of the mussel's life is still lacking (Moog *et al.*, 1998). Nevertheless, we discovered very recently that anomalies in nutritional conditions within mussel habitats as well as in their environments are clues to the species rarification (Hruska, 1999). According to this author, plant detritus entering the water from their surroundings serves as food source, and young mussels depend on this supply of detritus, which consists of small particles that penetrate the waters via underground transport processes.

2.5. **Threats and limiting factors**

The dramatic decline of *M. margaritifera* in Europe has run parallel to that of other naiads throughout the world. Bogan (1993) described the reasons for naiad rarification and extinction as habitat alteration or destruction, decline or extinction of fish hosts, commercial exploitation, and introduced species. Suitable habitat, i.e. Europe's clean salmonid rivers, is clearly disappearing.

The main reasons for the decline of *M. margaritifera* are anthropogenic influences on aquatic systems: eutrophication, impoundment, river regulation, drainage, sewage disposal, dredging, farms, new agricultural land use, loss of forests and natural river banks, water pollution, acidification and pesticides and introduced exotic fish species. Factors influencing fish host numbers are also responsible. However, data suggest that around 90% of European specimens disappeared in the twentieth century, overfishing for pearls being one of the main factors, particularly in Germany (Hessling, 1859; Boettger, 1954).

As we know that there is a very direct relationship between survivorship and availability of nutritive material from the environment, it seems that substances from areas subjected to intensive agriculture or forestry reach the water and could be directly related with the disappearance of the species (normally through the death of juveniles and consequent ageing of the populations).

The minimum population sub-unit is considered to be 500 reproductive individuals within 0.5 km of river (Irish data from Moorkens, 1999). This author summarises the threats to the species in her country as follows: 1- Nutrient enrichment. 2- Pollution incidents. 3- River bank erosion. 4- Forest plantation. 5- Road building. 6- Bog drainage and arterial drainage schemes. 7- Salmonid stocks. 8- Pearl fishing. 9- River modification. 10- Overgrazing. 11- Water abstraction. 12- Introduction of exotic species.

The spread of invasive species, such as the zebra mussel (*Dreissena polymorpha*), will be one of the main threats to native freshwater mussels, especially naiads.

2.5.1. *Habitat loss and habitat alteration*

The natural habitat of *M. margaritifera* was probably similar to those existing in the undisturbed rivers in the south of the Kola Peninsula. Although there are still some undisturbed rivers with *M. margaritifera* populations in a few countries of Northern Europe, this habitat is virtually being destroyed in most of the European countries where the species used to be common.

The main unwanted changes in freshwater or stream systems are mainly caused by the human activities listed below, either by direct physical destruction or due to sedimentation or eutrophication. For instance, a recruitment population of *M. margaritifera* numbering 5,000 individuals (the largest known population in Wales) was destroyed by dredging work on the Afon Ddu (Killeen, Oliver & Fowles, 1998).

Human activities responsible for freshwater pearl mussel decline through habitat alteration:

— Expansion and change in water bodies (river modifications, developments of drainage/abstraction schemes, flow regulation, hydro-electric schemes, fishery improvements) have been widely cited as unfavourable for freshwater mussels to thrive. One typical example is the construction of dams along freshwater mussel rivers that interrupt migratory trips of anadromous or catadromous fishes likely to harbour glochidia. Regarding river canalisation, there is a strong negative correlation between rivers that have been canalised and the presence of pearl mussels (Moorkens, 1996).

— Substrate mobilisation and dredging are directly responsible for freshwater mussel extraction (adults and juveniles) and secondary alterations to fish breeding areas.

— Deforestation causes a dramatic increase in river sedimentation. The disappearance of river shore trees leads to soil and nutrient losses through runoff and subsequent eutrophication and siltation.

— Eutrophication and decreasing water quality are also important factors working against freshwater mussel populations and river fish diversity. Although adult specimens can survive in eutrophicated waters, reproduction of those relict specimens in such conditions always presents a dangerous dependency. The juvenile stage is very susceptible to pollutants and a drop in water quality. Increasing organic material fills interstitial spaces in the substrate, which is the habitat of released juveniles.

— Bad agricultural land use causes alterations to river water quality (fertilisers, chemical pollutants, algal growth, eutrophication) and sedimentation systems. Cattle going into the river to drink cause river bank erosion and river bed siltation (Moorkens, 1999).

— Acid rains reduce salmonid populations and, therefore, freshwater pearl mussels, but it is not known to what extent acidification directly affects species (Cooksley, 1999).

2.5.2. *Human activities*

The abovementioned threats are of course due to human activities, but in this section we will deal with more direct human activities limiting species survivorship.

Fishing for the pearls of freshwater mussels has been known since the time of the Romans. More recently, data from Hessling (1859) give us an idea of the dangers derived from overfishing: between 1814 and 1857, 158,000 pearls were collected in the Bavaria region alone. This is a horrifying number bearing in mind that in order to find one pearl, nearly 3,000 mussels have to be opened (Hessling, 1859). In Ireland, Moorkens (1999) reports that at least five of the eight rivers containing breeding mussel populations were fished in the recent past.

Another threat directly associated with humans is the possible capture of specimens by poachers, once the risk to the species is already known. Collectors may be interested in having specimens of *M. margaritifera* due their value as pretty objects (pearls) and rare species.

Although not common now, log floating used to be a widespread technique on European rivers. In order to permit log floating, many rivers were altered by modifying the cross-section of their channels and, thus, the habitat of benthic fauna. This problem has been reported at least in Finnish rivers (Valovirta & Yrjänä, 1996).

The introduction of exotic fish species (i.e. rainbow trout *Oncorhynchus mykiss*) restricts the number of native fish as hosts of *M. margaritifera* glochidia.

The spread of invasive bivalve species favoured by Man (i.e. passive transport on the bottom of boats or in ballast water) reduces *M. margaritifera*'s ability to thrive. This has occurred in all countries invaded by the zebra mussel *Dreissena polymorpha*.

2.6. **Conservation status and current legal protection**

2.6.1. *International agreements*

— IUCN (International Union for Conservation of Nature and Natural Resources) 1996 Red Data List included the species as "Endangered" (IUCN, 1996).

— Habitats and Species Directive (European Union Directive on the Conservation of Natural Habitats and of Wild Flora and Fauna, -Directive 92/43/EEC). The species is listed in Annex II, which includes “animal and plant species of Community interest requiring the design of special conservation areas” and Annex V including “animal and plant species of Community interest whose taking in the wild and exploitation may be subject to management measures”. More recently, the EU adopted a new Directive (Directive 97/62/EEC) modifying the previous list. On this new list *M. durrovensis* has been listed as *Margaritifera durrovensis* (*Margaritifera margaritifera*), as well in Annexes II and V.

— Bern Convention (Council of Europe’s Convention on the Conservation of European Wildlife and Natural Habitats, 1979). The species was included in 1987 in Appendix III, “Protected fauna species”. However, until more scientific information on the taxonomic status of *M. durrovensis* is published, the Council of Europe does not plan to amend the list to include this taxon. The last meeting of the Group of Experts on Conservation of Invertebrates (Neuchatel, 2000) reviewed a draft version of this Action Plan (Araujo & Ramos, 2000) raising its quality and asked the Secretariat to prepare a reviewed version insisting in the need of international collaboration and a common protocol to standardize surveys of the species.

2.6.2. Listing of individual countries

- Austria

Although the species is endangered and threatened by extinction in most parts of its Austrian distribution (Moog *et al.*, 1998), it is not listed under nature protection laws. Nevertheless, it is protected by the fishery laws of Upper and Lower Austria. There are currently two projects aimed at protecting it, one in Upper Austria and the other in Lower Austria.

- Belgium

In addition to the abovementioned international agreements, the Bern Convention (Annexes III) and Habitats Directive (Annexes II and V), *M. margaritifera* is also included in the Regional Walloon Order of 3 June 1999 concerning the protection of several species of molluscs (M.B. 17/07/99).

- Czech Republic

On the River Blanice, where the main population lives, a species recovery project is currently in progress (Hruska, 1999). Nevertheless, problems from mechanical pollution due to the sand and other sediments during irregular flows and the absence of the necessary ecological agriculture in the river basin are threatening this population. Similar problems are affecting the other populations.

Margaritifera margaritifera appears on the Red Data List of Endangered and Rare Species of Plants and Animals of the CSFR 3-Invertebrates and among the critically endangered species on the Red List of Water Molluscs of the Czech Republic.

It is protected on a nationwide scale by Act No. 114/1992 Gazette on Protection of Nature and Landscape Conservation as a specially protected species in the “critically endangered” category. The River Blanice is a biogenetic reserve of the Council of Europe and River Teplá Vltava is part of a UNESCO biosphere reserve.

The conservation project is co-ordinated by the Nature Conservation and Landscape Protection Agency of the Czech Republic.

- Denmark

It was protected by Ministerial Order No. 67 of 4/2 1999 with later amendments, now under revision. The River Varde has been nominated as a protected site under the EC Habitats Directive because of the presence of *M. margaritifera*. The decline in the species is probably caused by past discharges of mercury into a lake belonging to the river system. At present, there are no plans to recover the species.

- Estonia

The species is protected on the Red Data List in Category 1 as an endangered species. According to the Act on Protected Natural Objects the species comes within Protection Category I. The site is also protected as habitat. The main threats affecting habitat are dredging, canalization and irregular water level.

- Finland

Finland's Red Data Book (1992) classifies *M. margaritifera* as a threatened species (threatened, vulnerable). The third edition of this Red Data Book (in preparation) will follow the new classifications of threatened species compiled by the IUCN. The species was protected by decree in Finland in 1955. More recently, the Habitats Directive and, in particular, the Natura 2000-network have been the key protection tools. In 1995 the Ministry of the Environment set a confiscation value of 3,500 FIM (585 EUROS) for a single *M. margaritifera*.

The World Wide Fund for Nature in Finland (WWF-SF), acting together with the Finnish Museum of Natural History, established (1978) a "*Margaritifera* working group" to inventory and study the distribution, ecology, morphology and protection of this species. Sub-aqua work is very prominent in the research of this group. During the last 22 years, the working group has checked freshwater pearl mussel populations in more than 70 rivers in different parts of Finland, and in some rivers in Russian Karelia and Estonia. All together, *Margaritifera* populations have been inventorised in more than 2000 km of one metre wide, underwater research transect.

In 1988, thirteen *M. margaritifera* sites were proposed as Natura 2000 sites (SCI). The on-going LIFE-Nature project (1997-2000) "Restoration of Fluvial Ecosystems Containing Pearl Mussels" includes the restoration of three riverine sites which have been proposed as SCI's for the Natura 2000 Network, as well as increasing public awareness. Seven organisations are taking part in this project, being the University of Helsinki the co-ordinator. The European Union granted 360,000 ECUs to start this LIFE Nature project. Recent progress in conserving *M. margaritifera* is summarised in the report of the conference "The freshwater pearl mussel in Europe: population status and conservation strategies" at Hof, Germany (Valovirta, pers. com.).

There are six main components in the project:

- 1) Inventory and history of *Margaritifera* populations. In the Korvuanjoki river (old pearl fishing river) only 500 specimens were found. The main reason was a thick layer of loose sand. Both of the southern rivers have been inventorised (SCUBA diving) very carefully by 5x5 m squares. In the river Pinsiö-Matalusjoki about 19,000 mussels were recorded and in the river Ruonanajoki 26,000 mussels.

The confiscation value (nature value) of these two populations is all together 150 millions FIM, which is about 2.5 million EUROS. This inventory has been comprehensive and the most accurate in Europe. It gave lot of new information about the optimum environment for *Margaritifera*. Moreover, the importance of the right water flow and the natural continuity of different river habitats, was quite a new result for the restoration of small rivers for freshwater pearl mussel, and for salmon (brown trout) too. It gave answers also to the question of why restoration work for salmon rivers has so often failed.

2) Restoration of river bed. For effective conservation of the freshwater pearl mussel two aspects of river quality must be addressed: The physical effects that contribute to river bed deterioration, and the parameters which define the quality of the water required by mussels in the river and drainage basin. During the last decade extensive projects have been started to restore the original habitat structure of canalised rivers in Finland. By 2000, more than 200 restoration projects had been completed, covering about 2000 separate rapid-flow sections of river channels. The right water flow and the natural continuity of different river habitats have proved very important in successful restoration of rivers for *Margaritifera* and for salmon, too. The Finnish "soft restoration technique" has been developed for *Margaritifera* rivers in the Natura-2000 areas. The core element of this technique is to restore rapid to rapid, stream to stream and still water to still water, using different construction methods, which re-establish the natural water dynamic specific to every river habitat. In restoring a seminatural river for *Margaritifera* we recognise the different types of river habitat and provide them with levels of natural water energy typical for every habitat type, as determined by channel slope and friction characteristics. It necessitates restoring what were originally rapids to rapids, streams to streams, still water to still water. In particular, restructuring a rapid into a series of short rapids and pools by deepening some sections and introducing bottom dams to others, will have a domino effect on other habitats both upstream and downstream, causing breakdown in their natural progression and consequent reduction in the sustainability of their conservation value.

3) Quality and flow of water. An extensive network of sampling stations for water analysis has been established on the watercourses since 1997, especially along the small southern rivers. The water quality parameters identified as active were to a significant extent established by comparisons made between sites with breeding mussel populations and sites with non-breeding populations or without any mussel population. The effects of water quality proved to be different for adult specimens and glochidial larvae. Continuous measurements of water flow, especially the overflow in spring and autumn, are suitable for recording in extreme habitat. Water volume, especially the volume of cool ground water, is important in southern Finland for the breeding of both *Margaritifera* and its brown trout host. For mussels, it is of paramount importance that, whatever restoration procedures are used, those procedures minimise the load of fine material entering the river water as a result of the restoration works, and leave the actual living area of the mussel untouched. The average suspended solids content we have recorded during the "soft" restoration process proved 6-15 times greater in the restoration areas than in control areas, but maximum quantities recorded hardly ever exceeded levels critical to the mussel populations. By contrast, where the "hard" restoration process was used the average suspended solids content was about 60 times greater than in the control areas, so that average quantities were way beyond the capacity of mussels to tolerate.

4) Artificial breeding of *Margaritifera* and transport of specimens. We have used artificial breeding in our project to help the glochidium of *Margaritifera* to find the gills of host fish in a laboratory and also under field conditions. The most critical epoch is one to two weeks in late summer or early autumn, when the mother mussel releases the larvae into the water of the river, and in the spring when the young mussels drop down into the river bottom. During these periods, good water quality is extre-

mely important for the lifecycle of *Margaritifera*. As a last conservation method we have used the translocation of freshwater pearl mussels from restoration areas. To do this we have adapted a method used for transporting salmon fry.

5) Host fish populations. We have used genetic tests to identify the original trout populations of the *Margaritifera* rivers. One product has been recognition that there are differences between brown trout populations from different rivers, in the degree of success with which they may be used to transplant mussel glochidia. Moreover, we now have information on which brown trout populations (from other rivers or fish-breeding farms), and which age classes, are the most suitable as a host fish for particular *Margaritifera* populations.

6) Publicity and nature conservation. Every year we are several times on TV and radio, we write numerous articles in newspapers and contribute lectures to national or international conferences. The project has a publication series of its own (Raakkuraportti = *Margaritifera* report). In 1999 the project had a display in the Wild Life Exhibition in Kuopio (>30000 visitors). We have used video films for documentation of the restored river sections. The Commission of the EU has selected this project as a "Success story" for the year 2000.

- France

Décret no. 99-615 du 7 juillet 1999. Annexe III: Protected Fauna Species. The species has received particular attention under the LIFE programme for the Allier and Loire basins.

An management plan has been approved in two regional parks (Morvan and Livradois-Forez) to avoid destruction of habitats suitable for the species.

- Germany

The species is listed as "threatened with extinction" on the "German Red List of Endangered Animals" (Jungbluth & Knorre 1998). It is strictly protected according to the German Federal Nature Conservation Law, off-take or changes in its habitat are generally prohibited. By region the situation is as follows:

— Niedersachsen: It is regarded as threatened with extinction. The main threat in this state is the expansion and change in water bodies. A large-scale conservation programme for the species (restoration of the natural river bank, removal of spruce forest, building of sandcatchers and extension of agricultural use) have led to juvenile recruitment. The programme costs 32,400,000 DM for the years 1989-2001 plus an additional 20,000 DM annually for direct species conservation measures (excluding salaries).

— Rheinland-Pfalz: It is listed as threatened with extinction on the Red List of Endangered Snails and Mussels in Rheinland-Pfalz (1994), but regarded as extinct in 1999. There is an intensive ongoing conservation project on the River Our in Luxembourg concerning as well the state of Rheinland-Pfalz due to the 'Condominium' status of this river.

— Hessen: It is listed as threatened with extinction on the Red List of Endangered Snails and Mussels of Hessen (1995).

— Bayern: It is listed as threatened with extinction on the last Red List of the Threatened Molluscs of Bavaria (1996) and strictly protected by the Fishery State Law (offtake is generally prohibited). A monitoring programme for the species in 17 selected streams was introduced by the Bavarian State Agency for Environmental Conservation including annual surveys of population status, water quality study, release of previously infested brown trout (since 1990) and breeding programmes under semi-natural conditions. A large-scale conservation programme to improve the habitat of the Zinnbach stream, near the Czech border, and other plans for freshwater and landscapes have been in progress since the mid-eighties. In 1995 a population of 5,000 individuals was translocated from an area threatened by road construction to an upstream area with an existing natural population. Monitoring in this case has not yet indicated the success or failure of the project. The project carried out in the Zinnbach (1985-1988), including construction of a separate waste-water collection channel and the establishment of protected areas on the direct stream bank, cost 1,165,000 DM.

— Thüringen: Listed as threatened with extinction on the last Red List of Endangered Mollusc Species of Thüringen (1993) and listed in the Fishery State Law of Thüringen, which includes the obligation to care and protect the species. There are no specific conservation measures with regard to *M. margaritifera*; however, there are general measures in operation in order to recover some parts of the Wettera stream. Nevertheless, every five years a population survey for large mussels is carried out in this state on behalf of the state Environment Agency in Thüringen.

— Sachsen: Listed as threatened with extinction on the last Red List of Endangered Terrestrial and Freshwater Molluscs of Sachsen (1996) and strictly protected by the Nature Conservation Act as well as the State Fishery Act (offtake or changes in habitat are generally prohibited). There are other state laws calling for the conservation of natural freshwater systems and ecological restoration following waste-water operations. The Wolfsbach is already protected because it is situated in the nature reserve NSG "Dreiländereck". Moreover, the Wolfsbach population and the one of the Raunerbach are proposed for being protected under the (FFH) Habitats Directive. In general, the aim to save the pearl mussel habitats in Sachsen was only partly achieved. Recent habitat changes have led to a dramatic decrease in populations.

- Ireland

In Northern Ireland legislation prohibits the sale of mussels (and pearls as their derivatives) (Schedule 7 of the Wildlife (NI) Order 1985). This but does not give the species protection from killing, injury or disturbance. Under Irish law, it is illegal to interfere with *M. margaritifera* (Statutory Instrument No. 112, 1990); thus pearl fishing is currently outlawed in the country (Moorkens, 1999). The hard-water form of pearl mussel *Margaritifera durrovensis* (Phillips, 1928) is restricted to one population in the River Nore, and enjoys the same level of legal protection under Irish and European law (Moorkens, 1999). A pamphlet alerting people to the conservation status of *M. margaritifera* has been produced as well as a recommendation to recognise the species *M. m. durrovensis* in the Bern Convention (Council of Europe, 1996). *M. m. durrovensis* is now listed separately as an Annex to *M. margaritifera* in the EU habitats Directive (van Helsdingen *et al.*, 1996). According to Costello *et al.*, (1996) the conservation of this species requires habitat restoration that will benefit the host fish and other species, *M. m. durrovensis* being a flagship species. Ongoing control of effluent discharges, river bank stabilisation and reduced phosphorous inputs from farmland have been proposed. In this report, the authors also proposed fencing using sponsors to finance stretches of the river. The co-ordination of all the tasks involved must be the responsibility of the nature conservation authorities (Costello *et al.*, 1996).

Recent estimations suggest that in one river of SW Ireland the population consists of over 2,000,000 reproductively mature individuals (Speight, 1998). This river's catchment is greatly affected by recent speculative forestry initiatives. Acquisition of large areas for conservation purposes have been suggested. The recent report that the zebra mussel (*Dreissena polymorpha*) has arrived in Ireland from Britain on pleasure boats is really alarming.

- Latvia

The species is included in Category 1 (endangered) of the Latvian Red Data Book (1985), new Latvian Red Data Book (1998) and Red Data Book of the Baltic Region. It is also included on the State List of Protected Animals (Latvian SSR Supreme Council Ruling 1957 and 1987) and List of Specially Protected Animals (Regulation of the Cabinet of Ministers in planning stage).

Forest zones along streams are protected through the prevention of felling. Part of one population is situated in Gauja National Park.

A project entitled "Inventories of Species and Habitats, Development of Management Plans and Capacity Building in Relation to Approximation of EU Birds and Habitats Directives" financed by Denmark's Nature Agency is currently in progress.

River regulation and general eutrophication over the last 50 years are reported to be the limiting factors. The threats are the expansion of successfully reintroduced beaver, dams, illegal wood cutting and overfishing of trout.

There is information on overfishing and export of pearl mussels to Sweden and Russia in the seventeenth and eighteenth centuries.

- Lithuania

Currently listed in the Red Data Book of Lithuania and protected by the Protected Animal, Plant, Fungi Species and Communities Act.

It is forbidden to destroy or damage protected species and their localities (breeding sites). The perpetrator has to compensate all damage and, if possible, restore the object to its former state (Augustasukas, 1998).

There are currently no plans or projects for the species.

- Luxembourg

Apart from the protection provided by the Bern Convention and the Habitats Directive, the species is included on the provisional Red List of Mollusc Species of Luxembourg promoted by the Lanis Office of Studies (Groh, 1997) in the category ***.

Margaritifera margaritifera is strictly protected by the "règlement grand-ducal" of 8th April 1986 concerning protection of certain species of wild fauna and as such under the legal status of a condominium.

As the Our is a frontier river, the protection provided by German legislation is also valid for the population in Luxembourg and *vice versa*. Legislation on pearl fishing (Perlrecht), species protection and nature protection laws (at Bund and at Länder) are also applicable.

The German area by the Our-basin containing an important population of *M. margaritifera* is already a protected area. Luxembourg's Nature Conservation Service is now pursuing the declaration of the equivalent area as protected. If this occurs, the pearl mussel could benefit from the synergy of the protection, management and development plans of both protected areas, which could be decisive for conservation of its habitat.

In order to protect fish hosts, the international commission on fishing has published special regulations for trout fishing in the River Our upstream of Vianden. Kayaking and river rafting boats have been forbidden between Trois Frontières and Dasbourg-Pont to protect the mussels from mechanical threats.

Since 1989 the "Direction des Eaux et Forêts de Luxembourg" has been carrying on a research programme on the physical, chemical and biological water conditions as well as on the threats. These data will be completed with parallel data from the German and Belgian governments.

Experiences with artificial trout infestations in captivity and subsequent release of these fish started in 1991 on the River Our following the Wellmann (1943) method. In 1998 and 1999 (seven to eight years after propagation) Jungbluth & Groh (in Groh, 1999) found juveniles of six to seven years old in the river.

Water quality has also been improved in most rivers in Luxembourg.

Monitoring of the population of *M. margaritifera* and its host fish in the River Our is ongoing since 1989.

A project on the Cartography of Molluscs has been underway since 1996 and involves drafting measures to protect the species.

Ideas for artificial raising of *M. margaritifera* under controlled conditions are under study as a joint project between German, Luxembourg and probably Belgian authorities.

- Norway

The species was given national protection on January 1st 1993 by the Act Governing Salmonids and Freshwater Fish (No. 47 of May 1992). This only protects the species against exploitation; however, the County Governor, as the local management authority, may issue permits if the proposed exploitation plan fulfils certain criteria. As far as is known, such exemptions have not yet been given. A number of nature protection areas (nature reserves) where the species occur have been established.

The national strategy to protect populations of the species involves mapping localities at national, regional (county) and local community levels. The information is fed into databases and marked on local maps to be used by area planners whenever exploitation plans are raised.

A recommendation for a ban on collecting and protection of the remaining habitats was recently concluded (Aagaard, 1998).

- Poland

Although the species is currently extinct, during the last few decades it was under strict legal protection, and now is still strictly protected by law.

- Portugal

No data available.

- Spain

Margaritifera margaritifera is protected under the following legislation:

— Transposition of the Bern Convention to Spanish legislation according to the General Provision of the Foreign Affairs Ministry of 26 May 1988 (BOE No. 136, of 07/06/1988, p. 17554), updated by the General Provision of the *Jefatura del Estado* of 13 May 1986 (BOE N° 235 of 01/10/1986): “Protected Wildlife Species”.

— Transposition of Directive 92/43/EEC to Spanish legislation by Royal Decree 1997/1995 of 7 December (Ministry of Agriculture, Fisheries and Food) (BOE No. 310, of 28 December, p. 37330). Annex II: “Animal and plant species of Community interest that require strict protection” and Annex V: “Animal and plant species of Community interest the collection of which in the wild and use may be the subject of management measures”. The modification of the Habitats Directive (Directive 97/62/EEC) was recently transposed to Spanish legislation by Royal Decree (1193/1998) of 12 June.

— National Endangered Species List. A project to include *M. margaritifera* on the National Red List as a ‘vulnerable’ species has already been approved by the National Fauna and Flora Committee. The species has not yet been included on any of the regional lists.

Two recent conventions between the Spanish Ministry for the Environment and the National Natural History Museum (Consejo Superior de Investigaciones Científicas -CSIC-) have been developing an updated bibliographical and distribution database (with field sampling) of the European threatened invertebrate species living in Spain, including *M. margaritifera*.

- Sweden

The species is included on the National Red List and has been classified as “vulnerable”. It is protected in the whole of Sweden by virtue of the fishery legislation.

At the Swedish Environmental Protection Agency, preparations are currently underway to produce a national action plan for *M. margaritifera*.

- United Kingdom

Freshwater pearl mussel is fully protected in Great Britain under Schedule 5 of the Wildlife and Countryside Act (1981 as amended 1998). In Northern Ireland legislation prohibits the sale of mus-

sels (and pearls as their derivatives) (Schedule 7 of the Wildlife (NI) Order 1985). This does not, however give the species protection from killing, injury or disturbance.

The species has been identified by the UK Government in its Biodiversity Action Plan (BAP) as a priority for conservation action (Biodiversity Steering Group 1995). Recently, the species has been the object of a national action plan drafted by Cooksley (1999) in consultation with the freshwater pearl mussel species action plan steering group (March 1999). Many organisations are involved in promoting and implementing the UK BAP. Scottish Natural Heritage (SNH) acts as the contact point and, in partnership with the Environment Agency act as lead partners.

37 Natura 2000 sites have been put forward, either specifically for *M. margaritifera* or have *M. margaritifera* present.

The objectives and targets of the national action plan are (Ball, 1996; Cooksley, 1999): 1- Establish the current status in UK and its ecological requirements at all stages of the life cycle. 2- Maintain and increase the size of existing populations. 3- Encourage re-colonisation. 4- Establish educational and monitoring programmes. 5- Determine effects of controlled exploitation in fished rivers and enforce legislation on pearl fishery practices.

Since 1994 intensive surveys for populations of the species have been carried out all over the country. Research into the ecological requirements of the species has been undertaken (postgraduate studies at Aberdeen University, the National Museum of Wales and Queen's University Belfast). A project entitled 'Culturing freshwater pearl mussel *Margaritifera margaritifera* in Northern Ireland. A first step towards the reintroduction of a threatened species' is currently underway. Specific freshwater mussel actions developed in the UK are summarised in Cooksley (1999).

3. GOALS AND OBJECTIVES

The main goal is, of course, effective conservation of *M. margaritifera* in each European country where populations exist, their habitat, future viability and long-term survival. Below is a list of tools to use and some points to be complied with in order to meet the proposed goals. Legislation, rational water use, research, management (including species re-introduction), monitoring and education/advisory work will be of the utmost importance. It is important to know that fertility of specimens is maintained in sparse populations and polluted rivers, indicating that they should recover if the causes of decline are removed (Bauer, 1988).

3.1 Policy and legislation

Due to the fact that *M. margaritifera* lives in fresh waters, conflict with people is of prime importance. Success is impossible if the corresponding administrations do not put all their co-ordinated efforts into the joint task. We know that the presence of a species name on a red list is not the final step towards its conservation, but only a necessary tool to work with. Once this tool is operating, the corresponding laws must be scrupulously adhered to. The items below summarise the proposed direct actions to be implemented to preserve the existing *M. margaritifera* populations.

To give the maximum level of protection to the species in all countries.

A positive dialogue and co-operation between Environmental, Hydraulic and Land-use administrations at national and regional levels is the primary objective and conservation of fresh waters inhabited by this threatened species one of the starting points.

New legislation regulating new water and land uses are needed. This should include: a) legislation to ensure water quality requirements for the species (when identified) form the basis for setting statutory water quality objectives, b) legislation to ensure that planning permission be sought for engineering work in rivers and channels where the species lives, and that such developments must be regulated, c) a guarantee that relevant management plans (e.g. agricultural, forestry, navigation) take account of the species interests and contribute to its conservation, d) ensure that local authorities are informed of *M. margaritifera* populations in their areas, the implications of the relevant legislation and contact points where advice may be sought.

National legislation to ensure that pearl fishing ceases in those countries with old traditions in this activity (e.g. Ireland, Germany) is also needed.

3.2. Species and habitat protection

All countries should designate areas and habitats of special protection for the species. Detailed conservation actions for each country are summarised below (Point 4 of this report). The protection of European freshwater systems harbouring populations of the species under the Natura 2000 Network and UNESCO biosphere reserves must be considered. Wider countryside measures need to be developed and implemented to protect the species, and its habitat, outside protected areas.

3.2.1. Develop and implement management plans to prevent further decline

A plan for rational use of water and land in the areas where the species lives must be designed. This can be carried out under major sustainable development projects in these areas.

It is a fact that irrigation farming around the world is under review. New social demands have given rise to generalised reflections to achieve sustainable ways of improving the quality and diversity of agricultural produce while attempting to conserve natural resources, which in this case are an endangered species and endangered freshwater habitats.

We cannot maintain, nor, of course, increase, existing levels of production and development without improving levels of productivity. This idea does not mean producing less, but rather handling the means of production in a controlled way, with minimal costs and without damaging the environment.

Under this general philosophy, specific commissions involving all political, scientific and socio-economic partners should be created to analyse the best conservation actions to be implemented in each case when new engineering or agricultural plans appear.

3.2.2. Conduct population surveys

Surveys of the rivers and water bodies harbouring the species or suspected to harbour it are still necessary in all countries. The results of these studies will be of great interest in the selection of the populations to be recovered in each European country. Although we already have current and realistic estimations of the populations in countries such as the Czech Republic, U. K. and Germany, there are other countries in which information is very scarce as for instance in Austria, Estonia, Lithuania, France, Spain or absent as in Portugal.

Rivers must be sampled using excavation and sieving techniques; relying solely on diver searches may lead to inaccurate assessment of population size structure. For assessing successful recruitment, total substratum removal of selected areas with subsequent sieving seems to be the best method (Miller & Payne, 1988; Richardson & Yokley, 1996).

3.3. Conflicts with humans

Only direct conflicts are going to be dealt with in this section due to the fact that major problems (water and land use, hydroelectric works, river regulation, etc.) have already been mentioned in this report.

Pearl fishing has historically been the most important activity acting against survivorship of the species and is still a real threat in some countries. Effective vigilance and a penalty policy can act in favour of the species.

The problem of collectors may be avoided with surveillance and information.

3.4. Public awareness

This is an important task that must be undertaken through joint action among scientists, local and national conservation authorities and other experts. Advisory publications, videos, conferences and other educational material dealing with the ecological importance of the species, legislative protection and the interest of its conservation for local people can be produced and distributed under an awareness campaign co-ordinated by Environmental Authorities and NGOs.

The items to be “sold” can be: water cleaning, beauty, uniqueness, natural interest, historical interest (pearl industry). In some cases, these items can be run under the heading of comprehensive rehabilitation of the habitats with social and economic advantages to local communities increasing their life quality.

The setting up of contact points in relevant areas should be encouraged not only to disseminate information, but also to target people who might be able to submit new *M. margaritifera* records.

Special attention should be given to civil servants responsible for surveillance in the relevant areas. They should receive suitable training on species identification and threat factors.

3.5. Population conservation

Indeed, a global conservation programme for the species, together with specific actions for the protection of each of the known populations of *M. margaritifera*, will probably be the key to the survival of the species.

Maintenance of all the different populations may be an essential source of genetic variability. This makes the survey to identify new populations very important. Special effort can be focused on aged populations because of the imminent risk of extinction.

Ways of conserving the populations will be dealt with in Section 4 of this report.

3.6. Artificial reproduction

The first step is to develop a programme to maintain breeding specimens of *M. margaritifera* in direct contact with salmon or trout in both monitored river sections and aquaria during the mussel's breeding season. Basic laboratories with aquaria and several raceways need to be available near the selected rivers in order to ensure that artificial infestation of fish occurs under the most natural conditions for the survival of newborn juveniles.

Successful results in artificial reproduction of *M. margaritifera* have recently been reported in the Czech Republic (Hruska, 1999).

3.7. Artificial propagation and reintroduction

Protected areas must be developed to house new generations of artificially bred mussels.

In this way, we have the opportunity to work with millions of recently released juveniles of *M. margaritifera* in the selected rivers of all countries. The juveniles can be hatched and kept in the selected river sections until they have grown several centimetres. These small mussels can be re-introduced in selected areas of the already known habitats in order to obtain reproductive mussels.

On the other hand, maintenance of recently released juveniles in the laboratory may be a good option. It will, however, require the study of different feeding methods and a suitable substratum. The advice of aquaculture specialists (marine clams and mussels) may be of prime interest in this regard.

3.8. Natural recovery

Probably the best method to recover the species is the re-introduction and maintenance of stocks of natural fish hosts in its former habitat. This possibility can only work in rivers where the fish can live and reproduce without problems.

Electro-fishing might be a good way of monitoring further infestation in the fish gills. Selected areas of the selected rivers could be used to investigate the success of mussel recruitment.

3.9. Monitoring and research

Further studies are needed to fill the gaps in knowledge of the known distribution of the species in Europe.

Research to investigate life history in different countries, genetic variation of the populations (for re-introduction and/or restocking experiences), substratum and water quality requirements, habitat requirements of juveniles or survival of re-introduced juveniles in natural ecosystems should be promoted.

The relation of mussel reproductive success to fish population levels has to be investigated in order to carry out a management plan of fish restocking, if necessary.

A very close follow-up of the infestation of fish, juvenile growth and their acclimatisation to selected habitats will be necessary (for at least five years) to evaluate the success of the recovery.

In the case of artificial propagation, averages of juvenile mortality have to be investigated in relation to suitable habitats.

A programme of regular monitoring of known populations also needs to be set up to identify further threats to the species.

4. CONSERVATION ACTIONS

It is difficult to provide a general list of measures to improve species conservation due to the individuality of each river or brook. General preservation measures must be developed after analysing each situation in detail (Moog *et al.*, 1998). Nevertheless, an evaluation of the data provided by the different countries with living populations of the species or from which it has recently been lost points to the same problem: anthropogenic influences on aquatic ecosystems. Given this fact, the priorities in the respective countries need to be established first. There is no sense in demanding a freshwater species be conserved at the same time as main roads, dams and macro-changes in the landscape make the freshwater ecosystem unsuitable for that species.

4.1. International actions

At present, the only known international initiative promoting the conservation of *M. margaritifera* is the current Council of Europe Action Plan.

Given that *M. margaritifera* only produces pearls in a minimum percentage of cases (one in every 3,000) and given the bad state of the European populations, as can be seen from the current report, it does not seem reasonable to maintain the species in Annex V of the Habitats Directive (animal and plant species of Community interest whose taking in the wild and exploitation may be subject to management measures). Listing *M. margaritifera* in Annex IV of the Habitats Directive (animal and plant species of Community interest that require strict protection) instead of in Annex V is highly encouraged.

It would be desirable for European freshwater ecosystems inhabited by *M. margaritifera* to be protected as Natura 2000 sites and/or UNESCO Biosphere Reserves.

A positive dialogue with the Russian authorities should be encouraged in order to gain mutual benefits on the basis of the huge populations currently thriving on the Kola Peninsula.

4.2. Listing of countries with necessary actions

- Austria

Reintroduction experiments using glochidia-infested fishes could take place in former pearl mussel habitats. Additionally, it is necessary to support the increase of remaining populations through fish hosts artificially infested with glochidia. Suitable measures to improve water quality in biological and chemical terms are an absolute necessity (Moog *et al.*, 1998).

- Belgium

Conservation measures have been proposed in a recent report on the pearl mussel situation in the River Our (Kinet & Libois, 1999) that could be the basis for a species action plan for all the rivers in the Walloon region. They include:

— The need for studies to obtain more information on the species distribution, population numbers and conservation status.

— To improve water quality by reducing the input of nitrates via a reduction of waste water mainly from communal areas and agriculture. It is necessary to identify the main sources of pollution, as well as solutions to avoid them within the framework of a general policy for water cleansing and to legislate or promote public awareness in order to avoid the use of fertilisers containing nitrates and phosphates in a 50-metres area along river banks. Complementary measures such as conservation of the already existing vegetation along river banks, bearing in mind that planting new trees and shrub-like vegetation can help to reduce the input of sediments. Economic compensation for owners of the surrounding private land should also be considered.

— Habitat restoration or protection. Besides other measures, it is considered to be very important to remove coniferous trees and replace them with shady riparian vegetation, particularly in stream sections with *M. margaritifera* and upstream of the watershed system. It is also important to reduce the input of suspended solids and fertilisers by stabilising river bank vegetation. Protection of areas that are important to the survival of the species could mean buying land in order to extend some of the existing protected areas.

— Maintain a healthy trout population.

— Undertake further assessment of known populations and initiate programmes of regular monitoring of the mussel population and of water quality.

— If water quality improves once the previous proposed measures have been adopted, a programme of infestation in captivity of young trout and subsequent release in areas containing rich populations of senescent mussels should be encouraged.

— In the Luxembourg section of the River Our, experiments in species restoration are currently in progress with infestation of captive trout and their subsequent release into the natural river ecosystems.

- Czech Republic

Establishing ecological agriculture in the Blanice river basin and maintaining the current quality of the water in the River Teplà Vltava are important measures to be continued. Other rivers containing freshwater pearl mussel populations must be optimised (agriculture and forest management) and cross-border co-operation improved in cases of borders with other countries (Austria and Germany).

Three kinds of species protection measures are proposed: a) short-term, consisting in the breeding of juvenile specimens from separate localities and subsequent monitoring, b) medium-term, saving the separate localities and stabilisation or decrease in the negative impact on habitats (acidification, chemical and mechanical pollution of rivers and brooks, use of chemical pollution in the economic exploitation of areas), c). long-term, reaching the required water quality and optimisation of the ecological

improvement of the economy in selected river basins. They will be the most important steps to achieve availability of the food spectrum needed for successful recruitment in mussel populations.

Given that the best populations are in protected areas far from human impact, the social and cultural influence of the proposed measures will be minimal. Nevertheless, the limiting factors in this task may be the absence of financial support.

- Estonia

There is a real need for an action plan for the species, but the question of financial resources has not been resolved. The proposed conservation measures involve the prevention of overheating of the water by planting trees along the river banks, keeping numbers of the mussel's temporary host glochidia high enough and working against river siltation.

- Finland

Short- and medium-term proposals to conserve the species and its habitats involve protection of the remaining populations, implementation of the Natura 2000 network, especially at sites where the pearl mussel is found, and continuation of the LIFE-project for the species.

The following could be considered as long-term proposals:

- establish freshwater pearl mussel reserves
- improve public awareness of pearl mussels
- carry out assessments on the effects of pollution on pearl mussels, in particular of suspended solid contents in water
- reduce the amount of suspended solid contents in water due to forest and peatland drainage, especially at the Natura 2000 sites
- carry out impact assessments prior to release of North American rainbow trout (*Oncorhynchus mykiss*)
- assess the need to establish breeding and reintroduction programmes
- promote co-ordination at national and international levels

The most significant socio-economic impact of the Natura 2000 network has been the uncertainty created by its effects on land use, water use and on local people's rights. The costs of implementing the network to areas including *M. margaritifera* are quite low due to the fact that the sites are either within existing nature conservation areas or protected by existing legislation (e.g. the Water Act). The required restoration measures may, in some cases, increase the implementation cost.

In order to help reproduce the species, infestations of brown trout with glochidia have been carried out in the laboratory. Indeed, salmon densities have been improved in areas with low numbers. Restocking mussels in different parts of the same river has had a success rate of over 90% after 16 years; this rate

falls to 50% when transfer of mussels from one river to another is involved. Information aimed at preventing further damage has been disseminated via the press, radio and TV.

- France

Surveys to know the actual species distribution and population numbers are urgent. Restricting river regulation and protecting freshwater systems containing populations of the species are the main actions needed in this country.

Species protection measures: protect the water systems inhabited by the species under the Natura 2000 Network. Consider the inclusion of the 10 better sites in the Ramsar Convention. To undertake agri-environmental measures at the Central Massif. Pedagogical information on the species and its ecological interest. Upbringing of the Conseil Supérieur de la Pêche guards to follow up the existing populations.

- Germany

— Niedersachsen: the ongoing decrease needs to be halted and the subsequent recovery of the population in the Lutter and Lachte streams promoted. Consequently, implementation of the sustainable development of landscape is indispensable in a large-scale conservation programme. Unless the input of sediments mobilised by human activities is halted, it will be impossible to ensure the survival of the pearl mussel.

Financial support will be needed in other stream systems in order to buy land.

Two research projects are currently being carried on in order to monitor the efficiency of the ongoing conservation measures in the Lutter watershed system: (i) Effects of disposal facilities for farmland drainage ditches which flow into the main streams of the Lutter watershed system. (ii) Sand freight monitoring in the Lutter watershed system and control of sand freight reduction due to the measures of the Lutter large-scale conservation project.

— Bayern: the ongoing conservation measures outlined in Section 2.6.2. will be continued in the forthcoming years. Both the monitoring programme and the conservation measures will be continued by the state authorities responsible for nature conservation and landscape planning.

The costs of the large-scale conservation project carried out in the Zinnbach area between 1985-1988, which included the construction of a separate wastewater collection channel and the “establishment of protected areas on the direct stream bank” amounted to a total of 1,165,000 DM.

— Sachsen: this state is currently preparing a broad species conservation programme involving co-ordination of different activities and measures within the management of agriculture, forestry, fisheries, water resources and traffic. It will also include crossborder co-ordination of conservation measures in Saxonia, Bavaria and the Czech Republic. Specific measures will be as follows:

- * Habitat conservation by protecting certain parts of the watershed system from the input of undesirable substances (nutrients and harmful substances through special treatment of waste waters from communal areas) and by reducing the input of these substances from

agricultural areas. To guarantee the water quality required for pearl mussel conservation. Both the stream areas of the watershed system upstream of the mussel population and large parts of the valley area will have to be included. To realise this aim, extensive areas will have to be purchased.

* Promotion of nature conservation concurring with agricultural management (extensification) in order to minimise substance inflow from streams in adjacent areas. Large parts of the valley and adjacent farmland as well as large areas upstream of the mussel populations have to be changed to extensive agricultural use (including changing arable fields to pasture land); the use of fertilisers and pesticides also has to be prohibited.

* Restriction on land use in areas with mussel populations. As mussels are very sensitive to the effects of different land uses in the catchment area of the watershed system, these uses have to be restricted or carried out in a way that avoids accidents. The different uses involve hydraulic engineering, large building projects, traffic and recreation.

* Measures to promote population development directly. These measures include concentrating extremely scattered mussel occurrences in appropriate stream sections by means of translocation for a certain time.

* Stabilisation of water quality and restoration of the conditions needed for the juvenile mussel development.

* Boosting mussel reproduction in streams by induced infestation of brown trout with glochidia and by halting the release of other fish species, especially rainbow trout.

* Direct relief action in the case of a meander breaking through or grass sod breaking off.

* Protection measures for bank vegetation.

* Measures to restructure and recover natural conditions in the stream area of the mussels as well as upstream areas of the watershed system.

* Develop an emergency action plan in the case of a disaster.

* Research activities in various aspects needed for successful conservation of the pearl mussel in the specific local area of Saxonia.

* Public relation activities. To achieve acceptance of the species conservation programme as a cross-department nature conservation programme and of the conservation measures, especially by local communities and land users, and to convey the message that the pearl mussel in Saxonia is part of the region's identity.

* Create the time-restricted post of co-ordinator for all the different conservation and promotional activities.

* Implement and finance the species conservation programme from various sources.

- Ireland
 - Establish reliable *M. margaritifera* survey methods (scuba and glass-bottomed bucket).
 - Salmonid host survey.
 - Check suitability of *M. margaritifera* habitat in target rivers.
 - Improve forestry management
 - Reduce siltation caused during forestry operations in conifer plantations. Install silt traps in conifer plantation drainage systems emptying into target rivers by agreement with forestry management.
 - Reduce influence of conifer plantation drainage waters. Block surface drains originating in conifer plantations, where these drains cross land with protected area status and empty into target rivers. Bottom-line 100m lengths of selected, additional conifer plantation surface-drains emptying into target rivers, using limestone chips or the equivalent.
 - Reduce influence of conifer plantations within the riparian zone.
 - Re-establish *M. margaritifera* populations (via transfer of glochidially-infested host fishes).
 - Improve river beds.
 - Contribute to the English-language dossier on *M. margaritifera* for installation on the Internet.
- Latvia

Establish protected areas - freshwater pearl mussel reserves. In this areas carry out: 1- Strict control of beavers, preventing the establishment of beaver dams. 2- Assess the hydrological and hydrobiological conditions at each pearl mussel location to determine the limiting factors. 3- Control the population of salmonid fishes. 4- Control of human impact (wood cutting, use of fertilizers, pollution, flow regulation, fishing).

Develop a monitoring programme on condition of oligotrophic waters and eutrophication. Apply methods of trout infestation with glochidia in salmonid breeding centre (nursery) Karli. Develop international protection for pearl mussel rivers crossing the borders of state. Continue survey of potential habitats for the freshwater pearl mussel in the Gauja, Salaca, Ogre, Aiviekste, Mēmele and Durbe River watersheds.

Two new recent problems have arisen: 1- financial resources for continuation of the Action Plan 2000 are not yet received and 2- the existence of a critically endangered population of pearl mussels in a private land (wood cutting, flow regulation, intensive pisciculture).

- Lithuania

Searching for new populations and protecting the respective habitats are essential measures.

- Luxembourg

It is important to continue research on the species's reproductive biology, demography, population structure, conservation status, situation of the host fish population and adult and juvenile habitats in the River Our.

In this area, measures to avoid negative environmental influences on the species should be taken by drawing up a plan to restore the areas that are susceptible to flooding, to restore the riverside vegetation, reduce the input of harmful substances, consolidate the river banks and restore optimal migration conditions for fishes by suppressing or transforming small river dams.

All catchments with potential populations of *M. margaritifera* should be identified in order to establish the real distribution of the species. In the case of new live populations being found, studies should be conducted on their size and age structure and on the situation of the host fish populations. If necessary, pilot reintroduction projects should be undertaken, which should include long-term monitoring programmes.

Fostering public awareness of the importance of the species and publicising that it is protected by law would help to conserve the species habitat.

Strict control of observance of the legislation.

Finally, a monitoring programme on the pearl mussel population needs to be established, including the results of the artificial propagation experiments and water quality monitoring.

- Norway

To increase knowledge about its biology and design a management strategy. Reduce erosion and secure areas with erosion. Prevention of clear-cutting, removal of riparian vegetation and other man-made alterations. Reduce the use of organic compounds, phosphorous and nitrogen. Introduction of adult mussels or infested fish, hatching and release of juveniles in suitable localities. In acidified areas, liming is important to secure remaining populations or create an environment for mussel reintroduction.

- Portugal

A survey of the rivers of the northernmost part of Portugal is needed, namely in the Montesinho and Peneda-Geres National Parks (Libois, pers. comm.).

- Spain

In a recent forum on the species, proposals were put forward for the urgent inclusion of *M. margaritifera* on the National Endangered Species List in order to have a legal tool to avoid new public works damaging the mussel populations. The proposed category is entitled 'sensitive to habitat alteration' or, provisionally, 'vulnerable' species.

Protect the populations in Galicia, Asturias and Castilla-León regions. To increase knowledge of the species biology and its ecological requirements throughout its life cycle. Inclusion of the species on the

Regional Endangered Species Lists and drafting of regional action plans. Avoid channelling and impoundment works in the main courses of the rivers and channels harbouring the species. To support the development of riparian forest and restoration of populations of autochthonous salmonids.

The rivers of the north-west corner of the Iberian Peninsula need to be surveyed for new populations.

- Sweden

Conservation measures will be formulated as part of the action plan currently being prepared. At present, liming is the most important single conservation measure being carried out. Future activities are likely to include: 1- protection of sites or river sections, where priority will be given to sites at which recruitment is functioning. 2- Improved site monitoring. 3- Information and advisory services to landowners, forest companies, fishermen. 4- Continued liming of waters, which is of benefit to the general environment.

- United Kingdom

Legislation was introduced in 1998 which gave *M. margaritifera* complete protection in England, Scotland and Wales. Action is needed to raise awareness of this legislation and to enforce prohibition of pearl fishing. Maintenance/improvement of water quality and protection from river engineering work must be addressed wherever populations occur. Every effort should be made to support measures that address the collapse of salmonid stocks and, where appropriate, to include freshwater pearl mussel interests in these measures.

A detailed programme of actions promoted by the national action plan to which lead agencies have agreed to contribute is summarised in Cooksley (1999).

A rivers project under EC LIFE funding is in progress. The project includes specific work for the pearl mussel, including investigations into water quality requirements.

5. REFERENCES

- Aagaard, K. 1998. Protection and management of invertebrates in Norway. In: *Convention on the conservation of European wildlife and natural habitats*. Group of Experts on Conservation of Invertebrates. Council of Europe. T-PVS (98) 18: 57.
- Adam, W. 1960. *Faune de Belgique. Mollusques*. Tome 1. *Mollusques terrestres et dulcicoles*. Institut Royal des Sciences Naturelles de Belgique. 402 pp. + 4 planches.
- Álvarez-Claudio, C., García-Rovés, P., Ocharan, R., Cabal, J. A., Ocharan, F. J. & Álvarez, M. A. A new record of the freshwater pearl mussel *Margaritifera margaritifera* L. (Bivalvia, Unionoida) from the River Narcea (Asturias, north-western Spain). *Aquatic Conservation: Marine and Freshwater Ecosystems* 10: 93-102.
- Araujo, R. & Ramos, M. A. 2000. Action Plan for *Margaritifera margaritifera* in Europe. *Council of Europe*, T-PVS (2000) 10. 38 pp.
- Araujo, R. & Ramos, M. A. In press. *Margaritifera margaritifera* (L.). In: *Invertebrados no insectos de la Directiva Hábitats en España*. Ramos, M. A. (Ed.). Serie Técnica del Ministerio de Medio Ambiente.
- Augustauskas, J. 1998. Report on the progress towards the conservation of Bern Convention Invertebrates in Lithuania, 1996-1998. In: *Convention on the conservation of European wildlife and natural habitats*. Group of Experts on Conservation of Invertebrates. Council of Europe. T-PVS (98) 18: 47.
- Ball, S. G. 1996. United Kingdom. In: *Convention on the conservation of European wildlife and natural habitats*. Group of Experts on Conservation of Invertebrates. Council of Europe. T-PVS (96) 33: 49-64.

- Bauer, G., 1986. The status of the Freshwater Pearl Mussel *Margaritifera margaritifera* L. in the South of its European Range. *Biological Conservation* 38: 1-9.
- Bauer, G. 1987. Reproductive strategy of the freshwater pearl mussel *Margaritifera margaritifera*. *Journal of Animal Ecology* 56: 691-704.
- Bauer, G. 1988. Threats of the freshwater pearl mussel in Central Europe. *Biological Conservation* 45: 239-253.
- Bauer, G., 1992. Variation in life span and size of the freshwater pearl mussel. *Journal of Animal Ecology* 61: 425-436.
- Biodiversity Steering Group. 1995. *Biodiversity: The UK Steering Group Report*. Vol. 2: *Action Plans*. HMSO, London.
- Birringer, D. & Truffner, U. 1990. *Studie der Ourbeckens zwischen Ouren und Dasburg im Hinblick auf die Schaffung eines Naturschutzgebietes*. Mém. Dipl. europ. Sc. Environ. F.U.L., Arlon. 189 pp + ann.
- Boettger, C. R. 1954. Flussperlmuschel und Perlenfischerei in der Lüneburger Heide. *Abhandlungen der Braunschweigischen Wissenschaftlichen Gesellschaft* 6: 1-40.
- Bogan, A. E. 1993. Freshwater bivalve extinctions (Mollusca: Unionoida): A search for causes. *American Zoologist* 33: 599-609.
- Cochet, G. 1999. Le statut des Margaritiferae de France (Mollusca: Bivalvia: Unionacea: Margaritiferae). *Vertigo* 6. 1996 (1999): 27-31.
- Cooksley, S. 1999. Work programme for the freshwater pearl mussel *Margaritifera margaritifera* UK species action plan 1999-2002. Draft.
- Costello, M. J., Moorkens, E., Larkin, M. & Kurz, I. 1996. The River Nore and its almost extinct pearl mussel: a priority for management and river restoration. In: *Convention on the conservation of European wildlife and natural habitats. Colloquium on Conservation, Management and Restoration of Habitats for Invertebrates: Enhancing Biological Diversity*. Killarney, Irlande, 26-29 May 1996. Council of Europe. T-PVS (96) 51: 31-37.
- Council of Europe. 1996. The present state of Bern Convention Invertebrates in Ireland. In: *Convention on the conservation of European wildlife and natural habitats*. Group of Experts on Conservation of Invertebrates. T-PVS (96) 33: 36.
- Chesney, H. C. G., Oliver, P. G. & Davis, G. M. 1993. *Margaritifera durrovensis* Phillips, 1928: Taxonomic status, ecology and conservation. *Journal of Conchology* 34(5): 267-299.
- Davis, G. & Fuller, S. 1981. Genetic relationships among recent Unionacea (Bivalvia) of North America. *Malacologia* 20: 217-253.
- Ferrant, V. 1892. *Beiträge zur Molluskenfauna des Grossherzogthums Luxemburg*. - Separat-Abdruck aus den: Mitteilungen der Fauna, Verein Luxemburger Naturfreunde, Jahrg. 1891-1892 (mit eigener Paginierung: 3-49); Luxemburg (P. Worré-Mortens).
- Ferrant, V. 1902. Faune des Mollusques terrestres et fluviatiles du Grand-Duché de Luxembourg. *Publications de l'Institut grand-ducal de Luxembourg, section des sciences naturelles et mathématiques*, 27(A): 1-232, 144 Abb. Luxembourg (M. Huss).
- Grande, C., Araujo, R. & Ramos, M. A., 2001. The gonads of *Margaritifera auricularia* (Spengler, 1793) and *Margaritifera margaritifera* (L, 1758) (Bivalvia: Unionoidea). *Journal of Molluscan Studies*, 67: 27-35.
- Groh, K. 1997. Liste der in Luxemburg vorkommenden und in den Nachbarländern auf der Roten Liste stehenden Weichtierarten mit einer Gefährdungs-Einschätzung für das Land Luxemburg. -2 S. Liste im Auftrag des Naturschutzbüros Lanius, Buschdorf (L) für das Naturschutzförderprogramm Luxemburg (unpublished); Hackenheim.
- Groh, K. 1999. Cahier espèce *Margaritifera margaritifera* (L. 1758). Grand-Duché de Luxembourg. Ministère de l'Environnement. 11 pp. Unpublished.
- Guilbot, R., Bellan-Santini, D., Deschamps-Cottin, M., Cochet, G., Descimon, H. & Dommaget, J. L. 1998. Examen des actions menées en France depuis 1996 pour la conservation des invertébrés inscrits à l'Annexe 2 de la Convention de Berne. In: *Convention on the conservation of European wildlife and natural habitats*. Group of Experts on Conservation of Invertebrates. Council of Europe. T-PVS (98) 18: 20-31.
- Hanstén, C, Pekkarinen, M. & Valovirta, I. 1997. Effect of transplantation on the gonad development of the freshwater pearl mussel, *Margaritifera margaritifera* (L.). *Boreal Environment research* 2: 247-256.

- Harms, W., 1907. Zur Biologie und Entwicklungsgeschichte der Flussperlmuschel (*Margaritana margaritifera* Dupuy). *Zoologischer Anzeiger* 31: 814-824.
- Harms, W., 1909. Postembryonale Entwicklungsgeschichte der Unioniden. *Zoologische Jahrbücher. Abteilung für Anatomie und Ontogenie der Tiere* 28: 325-386.
- Hessling, T. von, 1859. *Die Perlnmuscheln und ihre Perlen (Naturwissenschaftlich und geschichtlich mit Berücksichtigung der Perlgewässer Baerns)*. Leipzig. 372 pp.
- Hruska, J. 1999. Nahrungsansprüche der Flussperlmuschel und deren halbnatürliche Aufzucht in der Tschechischen Republik. *Heldia* 4(6): 69-79 + Tafel 2-5.
- INDUROT, 1999. *Informe preliminar sobre la situación de la madreperla de río (Margaritifera margaritifera) en España*. Universidad de Oviedo e Instituto de Recursos Naturales y Ordenación del Territorio, Ministerio de Medio Ambiente. Unpublished report. 12 pp.
- IUCN, 1996. *The 1996 IUCN Red lists of Threatened Animals*. IUCN. Gland, Switzerland and Cambridge, UK.
- Jungbluth, J. H. 1993. Analyses concernant la détérioration des eaux courantes dans les bassins hydrologiques de la Rur et du système Schwalm/Perlenbach. Annexe au point 3b de la réunion du 7 octobre 1993. 5 pp. Unpublished.
- Jungbluth, J. H., Coomans, H. E. & Grohs, H. 1985. Bibliographie der flussperlmuschel *Margaritifera margaritifera* (Linnaeus, 1758) (Mollusca: pelecypoda). *Instituut voor Taxonomische Zoologie (Zoologisch Museum). Universiteit van Amsterdam*, 41: 1-220.
- Jungbluth, J. H. & Knorre, D. V. 1998. Rote Liste der Binnenmollusken (Schnecken (Gastropoda) und Muscheln (Bivalvia)). In: Binot, M., Bless, R., Boye, P., Gruttke, H. & Pretscher, P. (Eds.). *Rote Liste gefährdeter Tiere Deutschlands. Schriftenreihe für Landschaftspflege und Naturschutz*, 55: 283-289.
- Killeen, I. J., Oliver, P. G. & Fowles, A. P. 1998. The loss of a freshwater pearl mussel (*Margaritifera margaritifera*) population in NW Wales. *Journal of Conchology Special Publication* 2: 245-250.
- Kinet, S. & Libois, R. 1999. Propositions pour la conservation des Unionides (Mollusques Bivalves) dans le bassin de l'Our (Parc Naturel Hautes-Fagnes Eifel). Université de Liège. Institut de Zoologie. Rapport Ministère Région Wallone, DGRNE, Service Conservation de la Nature, Namur, Octobre 1999. Unpublished.
- Larsen, B. M. 1997. Pearl mussel (*Margaritifera margaritifera* L.). A review of national and international knowledge. *NINA Fagrapport* 28: 1-51.
- Lydeard, Ch., M. Mulvey., and G. M. Davis. 1996. Molecular systematics and evolution of reproductive traits of North American freshwater unionacean mussels (Mollusca: Bivalvia) as inferred from 16S rRNA gene sequences. *Philosophical Transactions of the Royal Society of London Series B* 351:1593-1603.
- Machado, A. 1997. *Guidelines for action plans for animal species: planning recovery. Convention on the conservation of European wildlife and natural habitats (Bern Convention)*. Nature and Environment, No. 92. Council of Europe Publishing. 76 pp.
- Machordom, A., R. Araujo, D. Erpenbeck & M. A. Ramos. Conservation genetics of endangered Margaritiferidae (Bivalvia: Unionoidea). In prep.
- Meyers, T. R. & Milleman, R. E. 1977. Glochidiosis of salmonid fishes. I. Comparative susceptibility to experimental infection with *Margaritifera margaritifera* (L.). *Journal of Parasitology* 63: 728-733.
- Miller, A. C. & Payne, B. S. 1988. The need for quantitative sampling to characterize size demography and density of freshwater mussel communities. *American Malacological Bulletin* 6(1): 49-54.
- Moog, O., Ofenböck, T., Neseman, H. & Stundner, C. 1998. The freshwater pearl mussel *M. margaritifera* (L.) in Austria: Fundamental conservation measures for an endangered species. *Verhandlungen - Internationale Vereinigung für Angewandte Limnologie* 26: 2438-2443.
- Moorkens, E. A. 1996. *Studies on the biology and ecology of Margaritifera in Ireland*. Inpublished Ph. D Thesis, University of Dublin, Trinity College.
- Moorkens, E. A. 1999. *Conservation management of the freshwater pearl mussel Margaritifera margaritifera. Part 1: Biology of the species and its present situation in Ireland*. Irish Wildlife Manuals, No. 8. Dúchas, The Heritage Service, Dublin. 34 pp.
- Moorkens, E. A., and M. J. Costello. 1994. Imminent extenuation of the Nore freshwater pearl mussel *Margaritifera durrovensis* Phillips: a species unique to Ireland. *Aquatic Conservation: Marine and Freshwater Ecosystems* 4: 363-365.

- Nezlin, L. P., Cunjak, R. A., Zotin, A. A. y Ziuganov, V. V., 1994. Glochidium morphology of the freshwater pearl mussel (*Margaritifera margaritifera*) and glochidiosis of Atlantic salmon (*Salmo salar*): a study by scanning electron microscopy. *Canadian Journal of Zoology* 72: 15-21.
- Nobre, A. 1941. Fauna malacológica de Portugal II. Moluscos terrestres e fluviaes. *Memorias e Estudos do Museu de Zoologia da Universidade de Coimbra* 124: 1-278.
- Pekkarinen, M. y Valovirta, I., 1996. Anatomy of the glochidia of the freshwater pearl mussel, *Margaritifera margaritifera* (L.). *Archiv für Hydrobiologie* 137(3): 411-423.
- Phillips, R. A. 1928. On *Margaritifera durrovensis*, a new species of pearl mussel from Ireland. *Proceedings of the Malacological Society of London* 18:69-73.
- Ramos, M. A. 1998. Main progress towards the conservation of Bern Convention Invertebrates in Spain since 1996. In: *Convention on the conservation of European wildlife and natural habitats*. Group of Experts on Conservation of Invertebrates. Council of Europe. T-PVS (98) 18: 59-69.
- Richardson, T. D. & Yokley, P. 1996. A note on sampling technique and evidence of recruitment in freshwater mussels (Unionidae). *Archiv für Hydrobiologie* 137(1): 135-140.
- Ross, H. (1992). The reproductive biology of the freshwater pearl mussel *Margaritifera margaritifera* in Co Donegal. *Irish Naturalists' Journal* 24 (2): 44-50.
- Smith, D. G. & W. P. Wall. 1984. The Margariferidae reinstated: a replay to Davis and Fuller (1981), "Genetic relationships among recent Unionacea (Bivalvia) of North America". *Occasional Papers on Mollusks, Museum of Comparative Zoology (Harvard University)* 4(64):321-330.
- Speight, M. C. D. 1998. Progress report in Ireland, 1996-1998. In: *Convention on the conservation of European wildlife and natural habitats*. Group of Experts on Conservation of Invertebrates. Council of Europe. T-PVS (98) 18: 40-41.
- Stoltze, M. & Pihl, S. (eds.). 1998. *Roedliste 1997 over planter og dyr I Danmark*. Miljø-og Energiministeriet, Danmarks Miljøundersøgelser og Skov-og Naturstyrelsen.
- Van Helsdingen, P. J., Willemse, L. & Speight, M. C. D. 1996. *Background information on invertebrates of the Habitats Directive and the Bern Convention*. Part III. *Mollusca and Echinodermata*. Nature and environment, No. 81. Council of Europe: 399-529.
- Valovirta, I. 1996. Finland. Main progress in invertebrate conservation in Finland since 1994. In: *Convention on the conservation of European wildlife and natural habitats*. Group of Experts on Conservation of Invertebrates. Council of Europe. T-PVS (96) 33: 22.
- Valovirta, I. 1998. Finland. Progress in invertebrate conservation in Finland in 1996-1997. In: *Convention on the conservation of European wildlife and natural habitats*. Group of Experts on Conservation of Invertebrates. Council of Europe. T-PVS (98) 18: 18.
- Valovirta, I. & Yrjänä, T. 1996. Effects of restoration of salmon rivers on the mussel *Margaritifera margaritifera* (L.) in Finland. In: *Convention on the conservation of European wildlife and natural habitats*. Colloquium on Conservation, Management and Restoration of Habitats for Invertebrates: Enhancing Biological Diversity. Killarney, Irlande, 26-29 May 1996. Council of Europe. T-PVS (96) 51: 38-48.
- Vrignaud, S. 2000. *Raréfaction de la Moule perlière (Margaritifera margaritifera) dans l'Ardenne belge: bilan et perspectives (stage du 10/01 au 14/04/2000)*. Institut Supérieur Européen des Métiers de l'Environnement. TS + Conseiller Technique en Protection de l'espace Rural. Session 1999/2000. 41 pp + 10 Annexes.
- Wellmann, G. 1943. Fischinfektionen mit Glochidien der *Margaritana margaritifera*. *Zeitschrift für Fischerei* 41: 385-390, 5 Abb.
- Young, M. R. & Williams, J. C. 1984. The reproductive biology of the freshwater pearl mussel *Margaritifera margaritifera* (Linn.) in Scotland. II. Laboratory studies. *Archiv für Hydrobiologie* 100: 29-43.
- Ziuganov, V., A. Zotin., L. Nezlin., and V. Tretiakov. 1994. *The freshwater pearl mussels and their relationships with salmonid fish*. VNIRO Publishing House, Moscow, 104 pp.

6. ACKNOWLEDGEMENTS

We wish to thank E. Fernández-Galiano for the confidence he placed in the authors in asking them to prepare this report, which was drafted with the support of the Belgian Government's Ministry for the Walloon Region. We are also indebted to people and National Agencies listed in Appendix 2 that made possible the draft of this document. The previous draft

of this Action Plan received useful comments from the Group of Experts on Invertebrates (Neuchatel, 2000) and delegates of the different countries. Lesley Ashcroft checked the English version. Pepe Fernández kindly edited the text of this report.

7. APPENDIX 1: Summary on protection and management measures on *M. margaritifera*.

1. Water quality requirements of pearl mussel water bodies

Habitat requirements for the species are reported in point 2.4.2. of this document. Nevertheless here are summarized aspects of river quality for effective conservation based on Irish data (Moorkens, Valovirta & Speight, 2000):

- The water must be clean enough not to cause any direct stress on adult or juvenile mussels.
- Water quality must be high enough to ensure that eutrophication does not occur.

That means:

- Interstitial water chemistry must resemble free running water nutrient levels.
- Dissolved oxygen > 9 mg/l
- PH < 8 > 6.3
- B. O. D. < 3
- Total ammonia < 0.10 mg/l
- Conductivity < 200 µS/cm
- Nitrate < 1.7 mg/l
- Phosphates (ortho-P) < 0.06 mg/l
- Other parameters to ensure effective conservation and reproduction of the species are not yet evaluated.

2. Land and water use practices adjacent to pearl mussel habitats

It is known that chemical derivatives from agriculture uses and engineering works (dams, water derivations etc...) are among the main problems affecting successful thriving of *M. margaritifera* populations. Recommendations to avoid these problems can be:

- Ecological maintenance of water sources and small brooks of the rivers with populations of the species.
- Maintenance of natural forests and shore vegetation.
- Create buffer zones of 5-10 metres (minimum) width along rivers.
- To forbid gravel extraction, dam construction and water derivation in rivers to avoid siltation.
- To forbid all kind of farms along rivers to avoid eutrophication.
- To regulate agricultural fields along these rivers improving the soil quality (ecological maintenance of meadows by organic fertilization) in order to recuperate the healthy of the organic detritus in the water (base of the mussel food).

3. Methodological recommendations on artificial breeding and raising of young mussels.

- To develop protected areas with fertile populations of *M. margaritifera* to obtain gravid adults as source of glochidia.
- Installation of facilities (aquaria and raceways) to ensure controlled infestation of fish.
- To hatch the juveniles borned until several centimetres in both, indoor facilities and cages in the river in natural substrate of clean aerated gravel.
- Re-introduction of these juveniles in selected areas of the river.
- Re-introduction of infested host fish in selected areas of the river.

4. Methodological recommendations on the monitoring of known populations.

- Regular survey (one or two times yearly) of adult specimens by trained people (diving, snorkeling, viewing glass, etc...).
- Regular survey of suitable salmonid fishes (electro fishing) during glochidial release period.
- Regular survey of juvenile recruitment using excavation and sieving techniques in selected areas. (total substratum removal with subsequent sieving) .
- Regular monitoring of water quality (sources of pollution, flood changes, etc...).

8. APPENDIX 2: LIST OF REPORT CONTRIBUTORS

We would like to thank the Nature Agencies and Environment Departments of the respective countries and the people listed below for their help.

Austria: Harald Gross.

Belgium: Sylvain Vrignaud, Els Martens, Roland Libois and Pierre Gérard.

Czech Republic: Jan Plesnik, Handrij Härtel and Karel Absolon.

Denmark: Erik Buchwald and Claus Goldberg.

Estonia: Piret Kiristaja.

Finland: Ilkka Heikkinen, Jussi Soramäki and Ilmari Valovirta.

France: Gilbert Cochet.

Germany: Ute Grimm, Horst Gruttke and Christine Nagel.

Ireland: Martin Speight, Ferdia Marnell and Evelyn Moorkens.

Latvia: Mudite Rudzite.

Lithuania: Selemonas Paltanavicius.

Luxembourg: Marie-Paule Kremer.

Norway: Finn Kateras and Oystein Storkersen.

Poland: Zbigniew Glowacinski.

Portugal: Ana Isabel Queiroz, Luzia Sousa.

Spain: Carlos Nores, Consuelo Álvarez-Claudio and Pedro García-Rovés.

Sweden: Torsten Larsson, Lennart Henrikson.

U. K.: Deborah Procter & Martin Gaywood.



Figure 1. European countries with living population of *M. margaritifera*. Asterisk indicates the flourishing populations of the Kola Peninsula rivers.